



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$\therefore \frac{d^2x}{dt^2} \left(\frac{b^2+k^2}{b^2} \right) + \frac{gx}{R} = 0. \quad \therefore \frac{dx^2}{dt^2} + \frac{5gx}{7R} = 0.$$

$$\left(\frac{dx}{dt} \right)^2 = \frac{5g}{7R} (a^2 - x)^2 \quad \text{or} \quad T = \frac{\pi}{2} \sqrt{\frac{7R}{5g}}.$$

Let $R=20902410$ feet, $g=32.10614$ feet. Then $\sqrt{R/g}=806.871$.

$t=1267.433$ seconds $=21$ minutes, 7.433 seconds.

$T=1499.627$ seconds $=24$ minutes, 59.627 seconds.

t, T the time for the sphere to move from rest at the surface to the middle of the tunnel is constant for tunnels of all lengths and the same as for a tunnel passing through the center of the earth.

228. Proposed by J. E. ROSE, Mount Angel College, Mount Angel, Oregon.

AB, BC are two uniform rods freely hinged at B , whose weights are $W, 4W$, and lengths $2a, 4a$, respectively. The ends A, C of the rods are attached to small rings which slide on a rough horizontal wire. When the distance between the rings is the greatest for which equilibrium can exist, both of them are on the point of slipping. Find the coefficient of friction.

Solution by G. B. M. ZERR, A. M., Ph. D., Philadelphia, Pa., and S. LEFSEHETZ, Pittsburg, Pa.

Let R, R' be the reactions at C, A , respectively; $\mu R, \nu R'$ the friction at these points; $\angle BAC = \theta, \angle BCA = \phi$.

Then $R + R' = 5W, \mu R = \nu R' \dots (1, 2)$.

Taking moments around B we get

$$2R' \cos \theta = 2\nu R' \sin \theta + W \cos \theta \quad \text{or} \quad R' = W/[2(1 - \nu \tan \theta)] \dots (3).$$

$$2R \cos \phi = 2\mu R \sin \phi + 4W \cos \phi \quad \text{or} \quad R = 2W/(1 - \mu \tan \phi) \dots (4).$$

(1, 2) in (3) and (4) gives $9\mu - \nu = 10\mu \nu \tan \theta, 3\nu - 2\mu = 5\mu \nu \tan \phi$.

$$\therefore \frac{5}{\tan \phi + 6 \tan \phi} = \nu, \quad \frac{5}{4 \tan \theta + 9 \tan \phi} = \mu.$$

In this problem, $\mu = \nu$. $\therefore \mu = \frac{4}{5} \cot \theta = \frac{1}{5} \cot \phi$. $\therefore \tan \theta = 4 \tan \phi$.

Also $\sin \theta : \sin \phi = 2 : 1$, or $\sin \theta = 2 \sin \phi$.

$$\therefore \tan \phi = \frac{1}{2}. \quad \therefore \mu = \frac{2}{5}.$$